

MSSM confronts precision electroweak data and muon $g - 2$

D. Nomura (Tohoku U.)

特定領域研究「フレーバー物理の新展開」研究会 2012

Based on JHEP **11** (2011) 068

by **G.-C. Cho**, **K. Hagiwara**, **DN** and **Yu Matsumoto**



TOHOKU
UNIVERSITY

Plan of the talk

1. Introduction
2. Muon $g - 2$ vs MSSM
3. EW precision data vs MSSM
4. Summary




Introduction: Standard Model prediction for muon $g - 2$

QED contribution	$11\,658\,471.808 (0.015) \times 10^{-10}$	Kinoshita & Nio, Aoyama et al
EW contribution	$15.4 (0.2) \times 10^{-10}$	Czarnecki et al
Hadronic contribution		
LO hadronic	$694.9 (4.3) \times 10^{-10}$	HLMNT11
NLO hadronic	$-9.8 (0.1) \times 10^{-10}$	HLMNT11
light-by-light	$10.5 (2.6) \times 10^{-10}$	Prades, de Rafael & Vainshtein
Theory TOTAL	$11\,659\,182.8 (4.9) \times 10^{-10}$	
Experiment	$11\,659\,208.9 (6.3) \times 10^{-10}$	world avg
Exp – Theory	$26.1 (8.0) \times 10^{-10}$	3.3 σ discrepancy

(Numbers taken from HLMNT11, arXiv:1105.3149)

n.b.: hadronic contributions:



Full SM Result and Comparison with Other Groups




Fig. from HLMNT11, arXiv:1105.3149

Muon $g - 2$:

- ✓ Powerful probe for New Physics at TeV scale
- ✓ $\sim 3\sigma$ deviation between exp and theory (SM)
 \implies Signal of new physics?

Electroweak (EW) precision data:

- ✓ Useful probe for New Physics
- ✓ Some years ago the final LEP Z -pole data appeared ([hep-ex/0509008](https://arxiv.org/abs/hep-ex/0509008))

A natural question:


Suppose that the MSSM is responsible for the muon $g - 2$ anomaly. **Where is the SUSY parameter region favored by the final LEP EW data?** —

Important question to study BEFORE the LHC

SUSY Contributions to Muon $g - 2$





Suppose that the $\sim 3\sigma$ deviation is due to SUSY...

Leading SUSY contributions in the m_Z/m_{SUSY} expansion:



In most cases, the $\tilde{\chi}^\pm - \tilde{\nu}$ diagram (a) and/or the $\tilde{B} - \tilde{\mu}_{L/R}$ diagram (b) dominate. (Lopez-Nanopoulos-Wang, Chattopadhyay-Nath, Moroi, ...)


MSSM Contributions to Muon $g - 2$



x-axis: M_2
 (gaugino mass)

y-axis: $m_{\tilde{l}}$
 (slepton mass)

Muon $g - 2$ at MSSM sample points



No.	$\tan \beta$	μ	M_2	$m_{\tilde{E}}$	(a)	(b)	(c)	(d)	(e)	(a)-(e)	total
1	10	200	150	300	29.6	1.1	0.7	-2.9	-1.3	27.2	25.0
2	10	200	450	120	27.5	8.8	3.3	-7.1	-6.7	25.9	25.9
3	10	800	150	200	14.3	16.2	0.6	-2.7	-1.3	27.1	27.1
4	10	800	500	150	6.9	21.3	1.0	-2.5	-2.1	24.7	24.3
5	50	800	150	550	26.9	2.4	0.5	-2.6	-1.0	26.3	26.0
6	50	800	900	280	18.0	18.0	2.5	-5.9	-5.1	27.7	27.6

The chargino diagram (a) and/or the Bino-smuon $_{L,R}$ diagram (b) dominate in all the sample points.

Selected SUSY models and muon $g - 2$

Selected SUSY Models

	$\tan\beta$	μ	$m_{\tilde{\mu}_L}$	$m_{\tilde{\mu}_R}$	A_μ	M_1	M_2
SG 1 (mSUGRA, $\tan\beta = 10$)	10	396	181	116	-445	103	193
SG 2 (mSUGRA, high $\tan\beta$)	50	762	585	465	-145	277	510
GM 1 (Gauge Med., high $\tan\beta$)	42	504	441	214	25	181	339
GM 2 (Gauge Med., $\tan\beta \sim 10$)	15	300	257	120	-39	169	327
MM1 (Mirage Med., $\alpha > 0$)	10	430	188	255	-465	170	258
MM2 (Mirage Med., $\alpha < 0$)	10	-572	253	108	245	-99	-248
MM3 (Mirage Med., $M_2 < M_1$)	10	534	200	237	509	224	173

Muon $g - 2$ in the Selected SUSY Models

	(a)	(b)	(c)	(d)	(e)	(a)-(e)	total	pull
SG 1	25.7	21.5	1.5	-5.2	-5.4	38.1	37.6	1.2
SG 2	20.0	4.8	1.0	-3.4	-2.8	19.5	19.4	-1.0
GM1	34.6	11.7	1.4	-5.3	-9.2	33.2	33.0	0.7
GM2	27.1	10.6	1.6	-5.0	-9.0	25.3	24.8	-0.3
MM1	19.4	7.2	1.4	-4.5	-1.9	21.7	21.7	-0.7
MM2	13.2	18.8	0.7	-2.7	-4.2	25.8	24.7	-0.4
MM3	19.6	7.9	1.1	-3.8	-1.8	23.0	23.1	-0.5

LHC excluded region




Fig. originally from D. Grellscheid et al, arXiv:1111.3365

Introduction to EW Precision Study

- **LEP-I** ('89 - '95): The Z -boson properties studied in great detail using 17 millions of Z boson decays. (Final report appeared in 2005: hep-ex/0509008)
- To confront the EW precision data with theory, the **S, T, U parameters** are useful ([Peskin+Takeuchi](#)).

$$\gamma \text{---} \bullet \text{---} \gamma = i e^2 \Pi_{QQ} g^{\mu\nu} + \dots$$

$$Z \text{---} \bullet \text{---} \gamma = i \frac{e^2}{c s} (\Pi_{3Q} - s^2 \Pi_{QQ}) g^{\mu\nu} + \dots$$

$$Z \text{---} \bullet \text{---} Z = i \frac{e^2}{c^2 s^2} (\Pi_{33} - 2s^2 \Pi_{3Q} + s^4 \Pi_{QQ}) g^{\mu\nu} + \dots$$

$$W \text{---} \bullet \text{---} W = i \frac{e^2}{s^2} \Pi_{11} g^{\mu\nu} + \dots$$

$$\alpha S \equiv 4e^2 [\Pi'_{33}(0) - \Pi'_{3Q}(0)],$$

$$\alpha T \equiv \frac{e^2}{s^2 c^2 m_Z^2} [\Pi_{11}(0) - \Pi_{33}(0)],$$

$$\alpha U \equiv 4e^2 [\Pi'_{11}(0) - \Pi'_{33}(0)].$$

In this talk, we use an improved version, **S_Z, T_Z and M_W** ([Hagiwara+Haidt+Kim+Matsumoto, Cho+Hagiwara](#)).

S_Z and T_Z (1)

We define the 'bar charges' as

$$\begin{aligned}\bar{e}^2(q^2) &\equiv \hat{e}^2(\mu) \left[1 - \bar{\Pi}_{T,\gamma}^{\gamma\gamma}(q^2) \right], & \bar{s}^2(q^2) &\equiv \hat{s}^2(\mu) \left[1 + \frac{\hat{c}(\mu)}{\hat{s}(\mu)} \bar{\Pi}_{T,\gamma}^{\gamma Z}(q^2) \right], \\ \bar{g}_Z^2(q^2) &\equiv \hat{g}_Z^2(\mu) \left[1 - \bar{\Pi}_{T,Z}^{ZZ}(q^2) \right], & \bar{g}_W^2(q^2) &\equiv \hat{g}_W^2(\mu) \left[1 - \bar{\Pi}_{T,W}^{WW}(q^2) \right],\end{aligned}$$

where $\bar{\Pi}_{T,V}^{AB}(q^2) \equiv [\bar{\Pi}_T^{AB}(q^2) - \bar{\Pi}_T^{AB}(m_V^2)]/[q^2 - m_V^2]$ and the hat means the $\overline{\text{MS}}$ coupling. In terms of the bar charges, the S , T and U parameters can be written as

$$\begin{aligned}\frac{\bar{s}^2(m_Z^2)\bar{c}^2(m_Z^2)}{\bar{\alpha}(m_Z^2)} - \frac{4\pi}{\bar{g}_Z^2(0)} &= \frac{S}{4}, \\ \frac{\bar{s}^2(m_Z^2)}{\bar{\alpha}(m_Z^2)} - \frac{4\pi}{\bar{g}_W^2(0)} &= \frac{S + U}{4}, \\ 1 - \frac{\bar{g}_W^2(0)}{m_W^2} \frac{m_Z^2}{\bar{g}_Z^2(0)} &= \alpha T.\end{aligned}$$

S_Z and T_Z (2)

The S , T and U parameters:

$$\begin{aligned}\frac{\bar{s}^2(m_Z^2)\bar{c}^2(m_Z^2)}{\bar{\alpha}(m_Z^2)} - \frac{4\pi}{\bar{g}_Z^2(0)} &= \frac{S}{4}, \\ \frac{\bar{s}^2(m_Z^2)}{\bar{\alpha}(m_Z^2)} - \frac{4\pi}{\bar{g}_W^2(0)} &= \frac{S+U}{4}, \\ 1 - \frac{\bar{g}_W^2(0)}{m_W^2} \frac{m_Z^2}{\bar{g}_Z^2(0)} &= \alpha T.\end{aligned}$$

The last eq. can be written as

$$\frac{1}{\bar{g}_Z^2(0)} = \frac{1 - \alpha T + \bar{\delta}_G}{4\sqrt{2}G_F m_Z^2},$$

We are more interested in physics at the Z pole \implies Replace $\bar{g}_Z(0)$ with $\bar{g}_Z(m_Z^2)$: (S_Z and T_Z parameters)

S_Z - T_Z Plane Analysis

1. Calculate $\mathcal{O}_i^{\text{th}}(\Delta S_Z, \Delta T_Z, \dots)$, where \mathcal{O}_i are EW precision observables ($\Gamma_Z, \sigma_h^0, A_f, \dots$).


2. Construct the χ^2 function as

$$\chi^2 = \sum_{i,j} (\mathcal{O}_i^{\text{th}}(\Delta S_Z, \Delta T_Z, \dots) - \mathcal{O}_i^{\text{exp}}) \times (V^{-1})_{ij} (\mathcal{O}_j^{\text{th}}(\Delta S_Z, \Delta T_Z, \dots) - \mathcal{O}_j^{\text{exp}}),$$


where V is the covariance matrix,

$$V_{ij} = (\delta \mathcal{O}_i^{\text{exp}})(\delta \mathcal{O}_j^{\text{exp}}) \rho_{ij}.$$

3. Find the minimum of χ^2 with respect to ΔS_Z , ΔT_Z etc. Draw the contours $\chi^2 - \chi^2_{\text{min}} = \text{const}$ if necessary.



EW Precision Data vs MSSM, (I) S_Z - T_Z plane analysis




Using the final LEP EW precision data, we can give a constraint on MSSM contributions to S_Z and T_Z .

Our Results:

- ✓ The SM with $m_H \sim 100$ GeV gives a good description.
- ✓ In the MSSM, light sfermions tend to be disfavored.

EW Precision Data vs MSSM, (II) S_Z - T_Z plane analysis




Using the final LEP EW precision data, we can give a constraint on MSSM contributions to S_Z and T_Z .

Our Results:

✓ All the sample points are within or close to the $1\text{-}\sigma$ favored region.


Problem in Jet Asymmetry Data?



The value of the effective mixing angle \bar{s}^2 determined from leptonic asymmetry data and that from jet asym. data do not agree very well

⇒ **problem in jet asym. data (or in the analysis)?**

EW Precision Data vs MSSM, (IV) fit w/o jet asymm. data



If we do not use the jet asymmetry data, the favored region shifts to the left. (Negative ΔS_Z is favored.)

✓ Light sleptons are favored.

Summary

- We have studied the favored parameter region of MSSM using the results of the muon $g - 2$ and the EW precision data.
- From muon $g - 2$: when $\tan \beta = 10$, the slepton mass of a few hundred GeV is favored. When $\tan \beta = 50$, the sleptons as heavy as 1 TeV are allowed within $1-\sigma$.
- In well-studied models like mSUGRA and Gauge Med. there still is some parameter region favored from muon $g - 2$ and EW precision data at large $\tan \beta$ (LHC data already exclude light squarks/gluinos).
- If we leave out the jet asymmetry data, light sleptons become more favored, which is favored from muon $g - 2$ as well.